



## Tracking the elusive cytotoxic T cell response in pigs

**Jungersen, Gregers; Nielsen, Morten; Overgaard, Nana Haahr; Frøsig, Thomas Mørch; Sørensen, Maria Rathmann; Welner, Simon; Strube, Mikael Lenz; Pedersen, Lasse Eggers; Lund, Ole; Golde, William T.**

*Total number of authors:*  
11

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Jungersen, G., Nielsen, M., Overgaard, N. H., Frøsig, T. M., Sørensen, M. R., Welner, S., Strube, M. L., Pedersen, L. E., Lund, O., Golde, W. T., & Buus, S. (2016). *Tracking the elusive cytotoxic T cell response in pigs*. Abstract from 11th International Veterinary Immunology Symposium, Gold Coast, Australia.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Session:** Concurrent Workshop 2 - VIC/IUIS Toolkit

## Tracking the elusive cytotoxic T cell response in pigs

**Gregers Jungersen**<sup>1</sup>, Morten Nielsen<sup>2,3</sup>, Nana H Overgaard<sup>1</sup>, Thomas Frøsig<sup>1</sup>, Maria R Sørensen<sup>1</sup>, Simon Welner<sup>1</sup>, Mikael L Strube<sup>1</sup>, Lasse E Pedersen<sup>1</sup>, Ole Lund<sup>2</sup>, William T Golde<sup>4</sup>, Søren Buus<sup>5</sup>

1. *National Veterinary Institute, Technical University of Denmark, Frederiksberg, Denmark*
2. *Department of Systems Biology, Technical University of Denmark, Lyngby, Denmark*
3. *Instituto de Investigaciones Biotecnológicas, Universidad Nacional de San Martín, Buenos Aires, Argentina*
4. *Plum Island Animal Disease Center, United States Department of Agriculture, Orient Point, New York, USA*
5. *Department of Immunology and Microbiology, University of Copenhagen, Copenhagen, Denmark*

Quantitative and qualitative assessment of antigen-specific cytotoxic T cell (CTL) responses in pigs is not a straightforward process. Through the years we have developed a series of reagents, tools and protocols to characterize peptide-specific CTL responses in pigs.

The most common recombinant **SLA heavy chains were produced** and peptide binding motifs were determined by assays measuring the **affinity and stability of the peptide-SLA complex** (pSLA) interaction. These results have been used to **train neural networks** to predict the binding of any pSLA (<http://www.cbs.dtu.dk/services/>). Recombinant SLA molecules complexed with verified binding peptides can be assembled to **SLA multimers for staining of peptide-specific CTLs**, and measured by flow cytometry, as we have shown with **FMDV and influenza**. This, however, requires **SLA-matched pigs** for which we have developed two methods: a sequence-based, **high-resolution SLA genotyping** method by standard PCR for specific detection of eight in-house SLA molecules; and a **next-generation sequencing** method for parallel detection of up to 50 samples of barcoded cDNA PCR products spanning exon 2 and 3. The latter for a wider characterization of expressed alleles in candidate pigs.

The *in vivo* generation of CTL responses to antigens following peptide immunizations is thought to require **cross-presentation in appropriate dendritic cells (DC)**. In mice this was linked to targeting of CD103<sup>+</sup>DCs recruited after intraperitoneal immunizations. We have therefore developed a protocol for **intraperitoneal delivery of peptides** formulated in poly(I:C)/MMG-decorated liposomes (**CAF09**) to investigate the influence of peptide dose on the generation of CTL vs. antibody responses. Finally, the induced CTL killing was assessed by an ***in vivo* cytotoxicity assay**, where purified autologous PBMCs, fluorescently labeled and pulsed with target peptides, were reinjected into the donor. The *in vivo* killing of peptide-pulsed cells was measured by flow cytometry relative to non-pulsed PBMCs at different time points after cell transfer.